

the solution thus obtained in a suitable volumetric flask with sufficient 0.9 percent saline solution to obtain a stock solution containing about 2,500 micrograms of imipenem and 2,500 micrograms of cilastatin per milliliter. Transfer a 10-milliliter aliquot of this solution to a 50-milliliter volumetric flask and dilute to volume with phosphate buffer, 0.001M, to obtain a solution containing 500 micrograms of imipenem and 500 micrograms of cilastatin per milliliter (estimated).

(iii) *Calculations*—(A) *Imipenem and cilastatin potency*. Calculate the micrograms of imipenem and cilastatin per milligram as follows:

$$\frac{\text{Milligrams of imipenem or cilastatin}}{\text{per milligram}} = \frac{A_U \times P_S \times d}{A_S \times 1,000 \times W_S}$$

where:

A_U =Area of the imipenem or cilastatin peak in the chromatogram of the sample (at a retention time equal to that observed for the standard);

A_S =Area of the imipenem or cilastatin peak in the chromatogram of the imipenem or cilastatin working standard;

P_S =Anhydrous imipenem or cilastatin activity in the respective working standard solution in micrograms per milliliter;

d =Dilution factor of the 10 samples; and

W_S =Net contents of 10 containers in grams (gross weight of 10 containers in grams—tare weight of 10 containers in grams).

(B) *Imipenem and cilastatin content*. Calculate the imipenem or cilastatin content of the container as follows:

$$\frac{\text{Milligrams of imipenem or cilastatin per milligram}}{\text{milligram}} = \frac{A_U \times P_S \times d}{A_S \times 1,000}$$

where:

A_U =Area of the imipenem or cilastatin peak in the chromatogram of the sample (at a retention time equal to that observed for the standard);

A_S =Area of the imipenem or cilastatin peak in the chromatogram of the imipenem or cilastatin working standard;

P_S =Anhydrous imipenem or cilastatin activity in the imipenem or cilastatin working standard solution in micrograms per milliliter; and

d =Dilution factor of the sample.

(2) *Sterility*. Proceed as directed in § 436.20 of this chapter, using the method described in § 436.20(e)(1).

(3) *Pyrogens*. Proceed as directed in § 436.32(a) of this chapter, using a solution containing 5.0 milligrams of imipenem per milliliter except inject 10 milliliters per kilogram of rabbit weight.

(4) *Loss on drying*. Proceed as directed in § 436.200(a) of this chapter.

(5) *pH*. Proceed as directed in § 436.202 of this chapter.

[58 FR 26670, May 4, 1993]

§ 441.220b Imipenem monohydrate-cilastatin sodium for injection.

(a) *Requirements for certification*—(1) *Standards of identity, strength, quality, and purity*. Imipenem monohydrate-cilastatin sodium is a dry mixture of imipenem monohydrate, cilastatin sodium, and sodium bicarbonate packaged for dispensing. Its potency is satisfactory if it contains not less than 400 micrograms of imipenem and not less than 400 micrograms of cilastatin per milligram. Its imipenem content is satisfactory if it is not less than 90 percent and not more than 115 percent of the number of milligrams of imipenem that it is represented to contain. Its cilastatin content is satisfactory if it is not less than 90 percent and not more than 115 percent of the number of milligrams of cilastatin that it is represented to contain. It is sterile. It is nonpyrogenic. Its loss on drying is not more than 3.5 percent. When reconstituted as directed in the labeling, its pH is not less than 6.5 and not more than 8.5. The imipenem monohydrate used conforms to the standards prescribed by § 441.20a(a)(1).

(2) *Labeling*. It shall be labeled in accordance with the requirements of § 432.5 of this chapter.

(3) *Requests for certification; samples*. In addition to complying with the requirements of § 431.1 of this chapter, each such request shall contain:

(i) Results of tests and assays on:

(a) The imipenem monohydrate used in making the batch for potency, sterility, pyrogens, loss on drying, specific rotation, identity, and crystallinity.

(b) The batch for imipenem potency, cilastatin potency, imipenem content, cilastatin content, sterility, pyrogens, loss on drying, and pH.

(ii) Samples, if required by the Director, Center for Drug Evaluation and Research

(a) The imipenem used in making the batch: 10 packages, each containing approximately 500 milligrams.

(b) The batch:

(1) For all tests except sterility: A minimum of 20 immediate containers.

(2) For sterility testing: 20 immediate containers, collected at regular intervals throughout each filling operation.

(b) *Tests and methods of assay*—(1) *Imipenem and cilastatin potency and content*. Determine the potency of the sample in micrograms per milligram of both imipenem and cilastatin and the milligrams of both imipenem and cilastatin per container. Proceed as directed in § 441.20a(b)(1) of this chapter, preparing the cilastatin reference standard solution, the sample solution and calculating the imipenem and cilastatin potency and content as follows:

(i) *Cilastatin reference standard*. Accurately weigh approximately 25 milligrams of the cilastatin reference standard into a 50-milliliter volumetric flask. Immediately prior to analysis, add 10 milliliters of 0.9 percent saline solution and 1 milliliter of 0.1 percent bicarbonate solution. Add phosphate buffer, 0.001M, and shake until dissolved. Sonicate, if necessary, but for no longer than 1 minute. Dilute to volume with phosphate buffer, 0.001M, to obtain a solution containing approximately 500 micrograms of cilastatin per milliliter. Mix well and inject immediately.

(ii) *Preparation of sample solutions*—(a) *Imipenem and cilastatin potency (micrograms of imipenem and cilastatin per milligram)*. Remove the metal seal from each of 10 containers and determine gross weight in grams. Dissolve and wash out the entire contents of each container with 0.9 percent saline into an appropriate size volumetric flask to give a concentration of 5 milligrams per milliliter each of imipenem and cilastatin. Further dilute with phosphate buffer, 0.001 M, to obtain a solution containing 500 micrograms each of imipenem and cilastatin per milliliter (estimated). Wash each stopper and container with small quantities of acetone or methanol three

times being careful not to wet the container labeling. Allow the containers to air dry about 3 hours or to constant weight. Weigh each container and stopper to determine tare weight in grams.

(b) *Imipenem and cilastatin content (milligrams of imipenem and cilastatin per container)*. Reconstitute the sample as directed in the labeling, except use 0.9 percent saline solution as the reconstituting fluid. Then, using a suitable hypodermic needle and syringe, remove all of the withdrawable contents if it is represented as a single-dose container; or, if the labeling specifies the amount of potency in a given volume of the resultant preparation, remove an accurately measured representative portion from each container. Accurately dilute the solution thus obtained in a suitable volumetric flask with sufficient 0.9 percent saline solution to obtain a stock solution containing about 2,500 micrograms of imipenem and 2,500 micrograms of cilastatin per milliliter. Transfer a 10-milliliter aliquot of this solution to a 50-milliliter volumetric flask and dilute to volume with phosphate buffer, 0.001M, to obtain a solution containing 500 micrograms of imipenem and 500 micrograms of cilastatin per milliliter (estimated).

(iii) *Calculations*—(a) Calculate the micrograms of imipenem and cilastatin per milligram as follows:

$$\frac{\text{Micrograms of imipenem or cilastatin}}{\text{per milligram}} = \frac{A_u \times P_s \times d}{A_s \times 1,000 \times W_s}$$

where:

A_u =Area of the imipenem or cilastatin peak in the chromatogram of the sample (at a retention time equal to that observed for the standard);

A_s =Area of the imipenem or cilastatin peak in the chromatogram of the imipenem or cilastatin acid working standard;

P_s =Anhydrous imipenem or cilastatin activity in the respective working standards solutions in micrograms per milliliter;

d =Dilution factor for the 10 samples; and

W_s =Net contents of 10 containers in grams (gross weight of 10 containers in grams-tare weight of 10 containers in grams).

(b) Calculate the imipenem or cilastatin content of the container as follows:

$$\frac{\text{Milligrams of imipenem or cilastatin per container}}{A_u \times P_s \times d} = \frac{A_s \times P_s \times d}{A_s \times 1,000}$$

where:

A_u =Area of the imipenem or cilastatin peak in the chromatogram of the sample (at a retention time equal to that observed for the standard);

A_s =Area of the imipenem or cilastatin peak in the chromatogram of the imipenem or cilastatin working standard;

P_s =Anhydrous imipenem or cilastatin activity in the imipenem or cilastatin working standard solution in micrograms per milliliter; and

d =Dilution factor of the sample.

(2) *Sterility*. Proceed as directed in § 436.20 of this chapter, using the method described in paragraph (e)(1) of that section.

(3) *Pyrogens*. Proceed as directed in § 436.32(a) of this chapter, using a solution containing 5.0 milligrams of imipenem per milliliter except inject 10 milliliters per kilogram of rabbit weight.

(4) *Loss on drying*. Proceed as directed in § 436.200(a) of this chapter.

(5) *pH*. Proceed as directed in § 436.202 of this chapter.

[51 FR 11573, Apr. 4, 1986; 51 FR 22275, June 19, 1986, as amended at 55 FR 11582, Mar. 29, 1990. Redesignated at 58 FR 26669, May 4, 1993]

PART 442—CEPHA ANTIBIOTIC DRUGS

Subpart A—Bulk Drugs

Sec.

- 442.4 Cefaclor monohydrate.
- 442.6 Cefadroxil monohydrate.
- 442.7 Cefadroxil hemihydrate.
- 442.8a Sterile cefamandole nafate.
- 442.9a Sterile cefamandole sodium.
- 442.10 Cefazolin.
- 442.11a Sterile cefazolin sodium.
- 442.12 Cefoperazone sodium.
- 442.12a Sterile cefoperazone sodium.
- 442.13 Cefotaxime sodium.
- 442.13a Sterile cefotaxime sodium.
- 442.14 Cefoxitin sodium.
- 442.14a Sterile cefoxitin sodium.
- 442.15 Cefixime trihydrate.
- 442.16 Ceftazidime pentahydrate.
- 442.16a Sterile ceftazidime pentahydrate.
- 442.17 Ceftizoxime sodium.
- 442.17a Sterile ceftizoxime sodium.
- 442.18 Cefuroxime sodium.
- 442.18a Sterile cefuroxime sodium.
- 442.19 Cefuroxime axetil.

- 442.20a Sterile cefonicid sodium.
- 442.21 Cephaloglycin dihydrate.
- 442.22a Sterile cefmenoxime hydrochloride.
- 442.23a Sterile cephaloridine.
- 442.25a Sterile cephalothin sodium.
- 442.27 Cephalixin monohydrate.
- 442.28 Cephalixin hydrochloride monohydrate.
- 442.29a Sterile cephradine sodium.
- 442.40 Cephradine.
- 442.40a Sterile cephradine.
- 442.41 Cephradine dihydrate.
- 442.50a Sterile ceforanide.
- 442.52 Cefotetan.
- 442.53a Sterile cefotetan disodium.
- 442.54 Cefpodoxime proxetil.
- 442.55 Ceftriaxone sodium.
- 442.55a Sterile ceftriaxone sodium.
- 442.58a Sterile cefotiam dihydrochloride.
- 442.60 Cefpiramide.
- 442.69 Cefmetazole.
- 442.70a Sterile cefmetazole sodium.
- 442.80 Cefprozil.

Subpart B—Oral Dosage Forms

- 442.104 Cefaclor monohydrate oral dosage forms.
- 442.104a Cefaclor monohydrate capsules.
- 442.104b Cefaclor monohydrate for oral suspension.
- 442.106 Cefadroxil monohydrate oral dosage forms.
- 442.106a Cefadroxil monohydrate capsules.
- 442.106b Cefadroxil monohydrate tablets.
- 442.106c Cefadroxil monohydrate for oral suspension.
- 442.107 Cefadroxil hemihydrate oral dosage forms.
- 442.107a Cefadroxil hemihydrate capsules.
- 442.107b Cefadroxil hemihydrate tablets.
- 442.115 Cefixime trihydrate oral dosage forms.
- 442.115a Cefixime trihydrate for oral suspension.
- 442.115b Cefixime trihydrate tablets.
- 442.119 Cefuroxime axetil oral dosage forms.
- 442.119a Cefuroxime axetil tablets.
- 442.119b Cefuroxime axetil for oral suspension.
- 442.121 Cephaloglycin dihydrate oral dosage forms.
- 442.121a Cephaloglycin dihydrate capsules.
- 442.121b Cephaloglycin dihydrate for oral suspension.
- 442.127 Cephalixin monohydrate oral dosage forms.
- 442.127a Cephalixin monohydrate tablets.
- 442.127b Cephalixin monohydrate capsules.
- 442.127c Cephalixin monohydrate for oral suspension.
- 442.128 Cephalixin hydrochloride monohydrate tablets.
- 442.140a Cephradine for oral suspension.
- 442.140b Cephradine capsules.
- 442.140c Cephradine tablets.
- 442.141 Cephradine dihydrate capsules.